

# VERTOX

## CONTACT GEL

### A ready-to-use contact gel bait for the control of mice indoors. Contains brodifacoum 0.005% w/w and denatonium benzoate 0.001% w/w

For Professional use only for the protection of public health, stored products and materials. For use as a rodenticide only. NOT for amateur sale. Use biocides safely and sustainably. It is illegal to use this product for uses or in a manner other than that prescribed on this label.

#### Directions for use

Place contact gel where there are signs of mouse activity, like fresh droppings, inside or near nests, runways and mouse feeding places. Place contact gel points throughout the infested area.

Level of infestation	Bait points	Advice
Low	Place 5-15g per gel point every 5 metres	Place gel in bait boxes or covered/protected gel points to protect the gel from non-target animals. Contact gel can also be placed on strips of insulation tape or paper tape fixed to overhead pipe-ways and ductwork where non-target animals and humans do not have access to them. Contact gel can also be applied to holes in walls and partitions through which mice run. Regularly check contact gel consumption and replace consumed or spoilt contact gel until consumption has stopped. Repeat treatment in situations where there is evidence of new infestation (e.g. fresh tracks or droppings). It is important to ensure that gel strips cannot fall or become displaced onto sensitive areas during use.
High	Place 5-15g per gel point every 2 metres	

#### Instructions for safe handling and use

The ready-to-use gel bait is not intended to be added or mixed with any other product. A thorough survey of the infested site is essential, particularly in secluded and sheltered places, to determine the extent of the infestation. Product must be handled in a safe manner. For professional users the use of appropriate personal protective equipment (PPE) is advised. Wear suitable gloves. Do not decant product into unlabelled containers. Always read the label before use and follow the instructions provided. Avoid contact with the skin. Avoid all unnecessary exposure, in particular avoid ingestion. Bait must be securely deposited in baiting stations or other coverings so as to minimise the risk of consumption by other animals or children. Where possible, secure baits so that they cannot be dragged away. Dead rodent bodies, remains of unused bait or any fragments of bait found away from the bait station must be collected during all control operations to minimise the risk of consumption and poisoning to children, companion animals and other non-target animals. Remove all baits after treatment and dispose of them in accordance with local requirements. Unless under the supervision of a pest control operator or other competent person, do not use anticoagulant rodenticides as permanent baits. In

most cases, anticoagulant baits should have achieved control within 35 days. Should rodent activity continue beyond this time, the likely cause should be determined. When using do not eat, drink or smoke. Wash hands and face after application and use of the product, and before eating, drinking or smoking.

#### Public Area Use

When the product is being used in public areas and tamper-resistant bait stations are not used, the following must be implemented. When the product is being used in public areas, the areas treated must be marked during the treatment period and a notice explaining the risk of primary or secondary poisoning by the anticoagulant as well as indicating the first measures to be taken in case of poisoning must be made available alongside the baits. When tamper-resistant bait stations are used, they should be clearly marked to show that they contain rodenticides and that they should not be disturbed.

#### Additional safety information

Avoid release to the environment. Refer to special instructions/safety data sheet. To avoid risks to human health and the environment, comply with the instructions for use. Use bait containers clearly marked "poison" at all surface baiting points. Remove all remains of bait, dead rodents during and after treatment and dispose of safely. Prevent access to bait by children, domesticated animals and pets (particularly dogs, cats, pigs and poultry). It is illegal to use this product for the intentional poisoning of non-target, beneficial and protected animals. Harmful to wildlife.

#### Medical advice

In case of accident, suspected exposure or if you feel unwell, seek medical advice immediately (show the label where possible). In case of contact with skin, wash with soap and water. Remove and launder any contaminated clothing. In case of contact with eyes, remove contact lenses if present and rinse the eye slowly and gently with water for 15-20 minutes. Seek medical advice immediately, in case of ingestion or if swallowed seek medical advice immediately and show this container or label.

#### Note to Physician

Antidote – Vitamin K1 – under medical supervision. Effects result from the inhibition of blood coagulation, through interaction with vitamin K1. Poisoning symptoms may include: bruising easily, nose or gum bleeds, blood in stools or urine, excessive bleeding from minor cuts or abrasions. Note that poisoning symptoms may develop over the course of several days. UK medical professionals should contact the National Poisons Information Service ([www.npis.org](http://www.npis.org)) for further advice. In Eire, for information or to report a poisoning incident contact The National Poisons Information Centre, Beaumont Hospital, Dublin (01-8092166), retain the label for reference.

#### Resistance Information

The resistance status of the rodent population to brodifacoum should be taken into account when considering the choice of rodenticide to be used. Where resistance to brodifacoum has been shown or is suspected, resistant management strategies should be employed.

#### Precautions:

IF SWALLOWED: Immediately call a POISON CENTRE or doctor/physician. Store locked up in a closed container. Wear protective gloves. Keep out of reach of children. Keep / store away from food, drink and animal feeding stuffs. Do not eat, drink or smoke when using this product.

#### Directions for the safe disposal

This material and its container must be disposed of safely. In Eire dispose of packaging, remains of unused product and dead rodents in accordance with local regulations. If required, consult a professional waste operator or local authority. In the UK poisoned rodents may be disposed of by the waste producer at an incinerator, landfill or burial site permitted to accept that type of waste, or collected by a registered waste carrier and taken for disposal at a suitably permitted site. Remove all baits after treatment and dispose of them in accordance with local requirements. For further information on disposal contact the Environment Agency (<http://www.environment-agency.gov.uk>) or SEPA (<http://www.sepa.gov.uk>).

Batch No. See tag.

D.O.E. See tag.

PelGar supports:



International

VERTOX is a registered trade mark of PelGar International Ltd.  
Authorisation No. IE/BPA 70246 / UK-2014-0854

**PelGar International Ltd.**

Unit 13 Newman Lane, Alton, Hampshire, GU34 2QR.

Tel. +44 (0)1420 80744 **Web:** [www.pelgar.co.uk](http://www.pelgar.co.uk)

# 300g<sup>e</sup>

